



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06659 - Goat Anti-ACHE Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ACHE, YT, acetylcholinesterase (YT blood group), ARACHE, N-ACHE, OTTHUMP00000211347, OTTHUMP00000211349, OTTHUMP00000211356, acetylcholinesterase; apoptosis-related acetylcholinesterase

Official Symbol: ACHE

Accession Number(s): NP_000656.1; NP_001269378.1

Human GeneID(s): [43](#)

Important Comments: This antibody is expected to recognise isoform NP_000656.1 only (the ubiquitously expressed, hydrophilic form).

Immunogen

Peptide with sequence QFDHYSKQDRCSL, from the C Terminus of the protein sequence according to NP_000656.1; NP_001269378.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 70kDa band observed in HepG2 and Jurkat cell lysates (calculated MW of 67.8kDa according to NP_000656.1). Recommended concentration: 0.3-1µg/ml. Primary incubation 1 hour at room temperature.


Immunofluorescence: Strong expression of the protein seen in the cytoplasm, plasma membrane and nucleus of U2OS cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of HeLa cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Rat, Mouse




EB06659 (0.3 μ g/ml) staining of Jurkat (A) and (0.5 μ g/ml) HepG2 (B) cell lysate (35 μ g protein in RIPA buffer).
Detected by chemiluminescence.



EB06659 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton.

Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear, membrane and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB06659 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.