



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07079 - Goat Anti-Adenosine A2a Receptor Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ADORA2A, adenosine A2a receptor, HGNC:263, ADORA2, RDC8, hA2aR, adenosine A2 receptor, adenosine receptor subtype A2a

Official Symbol: ADORA2A

Accession Number(s): NP_000666.2

Human GeneID(s): [135](#)

Immunogen

Peptide with sequence HELKGVCPPEPPGLDD, from the C Terminus of the protein sequence according to NP_000666.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: This antibody has been successfully used in Western blot on Macaque: van der Putten C et al. (2009) PMID: 19494284. This antibody has been successfully used in Western blot on Mouse: He X et al. (2013) PMID: 23712033, and Zhao L et al. (2015) PMID: 26171727.

Immunofluorescence: Strong expression of the protein seen in the cytoplasm of HepG2 cells. Recommended concentration: 10µg/ml.

Species Reactivity

Tested: Human, Mouse, Macaque

Expected from sequence similarity: Human

Specific References

This antibody has been successfully used in WB on Mouse:

Zhao L, Liu YW, Yang T, Gan L, Yang N, Dai SS, He F.

The mutual regulation between miR-214 and A2AR signaling plays an important role in inflammatory response.

Cell Signal. 2015 Oct;27(10):2026-34.

PMID: 26171727

This antibody has been successfully used in the WB on Mouse:

He X, Hu JL, Li J, Zhao L, Zhang Y, Zeng YJ, Dai SS, He FT.

A feedback loop in PPARγ-adenosine A2A receptor signaling inhibits inflammation and attenuates lung damages in a mouse model of LPS-induced acute lung injury.


Cell Signal. 2013 Sep;25(9):1913-23.

PMID: 23712033

This antibody has been successfully used in WB on Macaque:

van der Putten C, Zuiderwijk-Sick EA, van Straalen L, de Geus ED, Boven LA, Kondova I,

IJzerman AP, Bajramovic JJ.
Differential expression of adenosine A3 receptors controls adenosine A2A
receptor-mediated inhibition of TLR responses in microglia.
J Immunol. 2009 Jun 15;182(12):7603-12.
PMID: 19494284



EB07079 Immunofluorescence analysis of paraformaldehyde fixed HepG2 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).