

## **UK Office**

### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

#### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$ 

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08913 - Goat Anti-APOBEC1 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** APOBEC1, apolipoprotein B mRNA editing enzyme, catalytic polypeptide 1, APOBEC-1, BEDP, CDAR1, HEPR, apolipoprotein B mRNA editing

enzyme, apolipoprotein B mRNA editing enzyme complex-1

Official Symbol: APOBEC1

Accession Number(s): NP\_001635.2

Human GeneID(s): 339

Non-Human GenelD(s): 11810 (mouse)

# **Immunogen**

Peptide with sequence DVFYDPRELRKEAC, from the internal region (near N Terminus) of the protein sequence according to NP\_001635.2.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx. 28kDa band observed in Mouse Spleen lysates (calculated MW of

28.2kDa according to Human NP\_001635.2 and 27.5kDa according to Mouse

NP\_112436.1). Recommended concentration: 1-3µg/ml.

# **Species Reactivity**

Tested: Mouse

Expected from sequence similarity: Human, Mouse



EB08913 ( $1\mu g/ml$ ) staining of Mouse Spleen lysate ( $35\mu g$  protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.