



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB10027 - Goat Anti-POU3F3 / BRN1 / OCT8 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: Brain-1, BRN1, OTF8, POU class 3 homeobox 3, POU domain, class 3, transcription factor 3, POU3F3, OTF-8, brain-specific homeobox/POU domain protein 1, brn-1, oct-8, octamer-binding protein 8, octamer-binding transcription factor 8

Official Symbol: POU3F3

Accession Number(s): NP_006227.1

Human GeneID(s): [5455](#)

Non-Human GeneID(s): 18993 (mouse), 192109 (rat)

Immunogen

Peptide with sequence C-HMLSHAHQWVTAL, from the internal region of the protein sequence according to NP_006227.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:4000.

Western blot: Approx 48kDa band observed in Mouse Spinal Cord lysates and approx. 50kDa in Mouse Brain lysates (calculated MW of 50.3kDa according to Human NP_006227.1 and 50.2kDa according to Mouse NP_032926.2). Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature.


Immunofluorescence: Strong expression of the protein seen in the nuclei of Neuro-2a cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of Neuro-2a cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Mouse

Expected from sequence similarity: Human, Mouse, Rat




EB10027 (1 μ g/ml) staining of Mouse Spinal Cord (A) and Brain (B) lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB10027 Immunofluorescence analysis of paraformaldehyde fixed Neuro-2a cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (2 μ g/ml), showing

nuclear staining. Actin filaments were stained with phalloidin (red) and the nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB10027 Flow cytometric analysis of paraformaldehyde fixed Neuro-2a cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.