



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09792 - Goat Anti-SNX1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: HsT17379, MGC8664, SNX1A, sorting nexin 1, sorting nexin 1A, Vps5, SNX1

Official Symbol: SNX1

Accession Number(s): NP_003090.2; NP_690039.1

Human GeneID(s): [6642](#)

Non-Human GeneID(s): 56440 (mouse), 84471 (rat)

Important Comments: This antibody is expected to recognize isoform a and c (NP_003090.2; NP_690039.1).

Immunogen

Peptide with sequence C-TNSSKPQPTYEELE, from the internal region of the protein sequence according to NP_003090.2; NP_690039.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 75kDa band observed in lysates of cell lines HeLa and U2OS (calculated MW of 59.1kDa according to NP_003090.2). This molecular weight is routinely observed by other sources. Recommended concentration: 0.1-1µg/ml. Primary incubation 1 hour at room temperature.

IHC: In paraffin embedded Human Uterus shows vesicular staining in the cytoplasm of endometrial epithelium. Recommended concentration, 3-5µg/ml.

Immunocytochemistry: Vesicular staining of SNX1 (green) in formaldehyde-fixed SH5Y5Y. Recommended concentration, 3-5µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB09792 (1 μ g/ml) staining of HeLa (A) and (0.1 μ g/ml) U2OS (B) cell lysate (35 μ g protein in RIPA buffer).. Detected by chemiluminescence.



EB09792 (3.8 μ g/ml) staining of paraffin embedded Human Uterus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB09792 (5ug/ml) staining of formaldehyde-fixed SH5Y5Y and detected with Alexa 488 in confocal microscopy.
Data obtained from Dr. Schallburg Nielsen, Aarhus University Denmark.