



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07652 - Goat Anti-ZDHHC8 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ZDHHC8, zinc finger, DHHC-type containing 8, ZDHHCL1, ZNF378, membrane-associated DHHC8 zinc finger protein, zinc finger, DHHC domain containing 8, zinc finger, DHHC domain like containing 1

Official Symbol: ZDHHC8

Accession Number(s): NP_037505.1

Human GeneID(s): [29801](#)

Non-Human GeneID(s): 27801 (mouse), 303796 (rat)

Important Comments: This antibody is expected to recognise isoform 2 (NP_037505.1) only.

Immunogen

Peptide with sequence C-QRDHPQLKTPPSK, from the internal region of the protein sequence according to NP_037505.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 80-85kDa band observed in Human Brain (Frontal Cortex) lysates (calculated MW of 81.4kDa according to NP_037505.1). Recommended concentration: 1-3µg/ml. An additional band of 48kDa was consistently observed, however this band was not blocked by the immunizing peptide and it is therefore a non-specific signal. We call for caution when used for other assays than Western blot.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig

Specific Reference

This antibody has been successfully used in Western blot on Mouse:

Avishek Mondal, Abhilash P. Appu, Tamal Sadhukhan, Maria B. Bagh, Rafael M. Previde, Sriparna Sadhukhan, Stanko Stojilkovic, Aiyi Liu, Anil B. Mukherjee

Ppt1-deficiency dysregulates lysosomal Ca⁺⁺ - homeostasis contributing to pathogenesis in a mouse model of CLN1 disease

J Inherit Metab Dis. 2022 Feb 12. doi: 10.1002/jimd.12485.

PMID: 35150145



EB07652 (0.1µg/ml) staining of Human Brain (Frontal Cortex) lysate (35µg protein in RIPA buffer) with (B) and without (A) blocking with the immunising peptide. Primary incubation was 1 hour. Detected by chemiluminescence.